

Jet Quenching with Detailed Balance in a Thermal QCD Medium *

Enke Wang^a and Xin-Nian Wang^{ba}

^aInstitute of Particle Physics, Huazhong Normal University, Wuhan 430079, China

^bNuclear Science Division, Lawrence Berkeley Laboratory, Berkeley, California 94720

The effect of detailed balance in induced gluon radiation for an energetic parton propagating inside a quark-gluon plasma is discussed. The stimulated thermal gluon emission reduces while absorption increases the parton's energy. The net contribution results in a reduction of the effective parton energy loss. The relative reduction of the parton energy loss is found to be important for intermediate parton energies and negligible for very high energies. This will increase the energy dependence of the parton energy loss and will affect the shape of suppression of moderately high p_T hadron spectra.

1. INTRODUCTION

Jet quenching in high-energy heavy-ion collisions suppresses large transverse momentum hadrons as compared to pp collisions at the same energy [1,2]. It provides a direct probe of the dense matter produced in high-energy heavy-ion collisions. Recent theoretical studies [3–7] of jet quenching have concentrated on gluon radiation induced by multiple scattering in medium. Because of the presence of thermal gluons in the hot medium the stimulated gluon emission and absorption by the propagating parton should also be taken into account consistently. Such a detailed balance is crucial for parton thermalization and should also be important for calculating the energy loss of an energetic parton in quark-gluon plasma (QGP)[8]. We will show that the partial cancellation between stimulated emission and thermal absorption results in a net reduction of parton energy loss induced by multiple rescattering. Though such a reduction is negligible for asymptotically large parton energy, it is still important for intermediate values of energy.

2. FINAL-STATE AND RESCATTERING-INDUCED ABSORPTION

In an axial gauge and in the leading-log approximation, the final-state radiation amplitude off a quark can be factorized from the hard scattering. Taking into account of both stimulated emission and thermal absorption in a thermal medium with finite temperature T , the probability of gluon radiation with energy ω can be written as

$$\frac{dP^{(0)}}{d\omega} = \frac{\alpha_s C_F}{2\pi} \int \frac{dz}{z} \int \frac{d\mathbf{k}_\perp^2}{\mathbf{k}_\perp^2} \left[N_g(zE) \delta(\omega + zE) + (1 + N_g(zE)) \delta(\omega - zE) \theta(1 - z) \right] P\left(\frac{\omega}{E}\right), (1)$$

*This work was supported by the NSFC under projects Nos. 19928511 and 10135030, and by the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

where $N_g(|\mathbf{k}|) = 1/[\exp(|\mathbf{k}|/T) - 1]$ is the thermal gluon distribution, C_F is the Casimir of the quark jet in the fundamental representation and the splitting function $P_{gq}(z) \equiv P(z)/z = [1 + (1 - z)^2]/z$ for $q \rightarrow gq$. In Eq.(1) the first term is from thermal absorption and the second term from gluon emission with the Bose-Einstein enhancement factor.

To define the effective parton energy loss, we only consider gluon radiation outside a cone with $|\mathbf{k}_\perp| > \mu$. Here μ should be a nonperturbative scale for parton fragmentation in the vacuum. In a hot QGP medium, it can be replaced by the gluon's Debye screening mass. Assuming the scale of the hard scattering as $Q^2 = 4E^2$, one has the kinematic limits of the gluon's transverse momentum, $\mu^2 \leq \mathbf{k}_{\perp max}^2 \leq 4|\omega|(E - \omega)$.

Subtracting the gluon radiation spectrum in the vacuum, one then obtains the energy loss due to final-state thermal absorption and stimulated emission. For asymptotically large parton energy, $E \gg T$, we have

$$\begin{aligned} \frac{\Delta E_{abs}^{(0)}}{E} &= \int d\omega \frac{\omega}{E} \left(\frac{dP^{(0)}}{d\omega} - \frac{dP^{(0)}}{d\omega} \Big|_{T=0} \right) \\ &\approx -\frac{\pi\alpha_s C_F}{3} \frac{T^2}{E^2} \left[\ln \frac{4ET}{\mu^2} + 2 - \gamma_E + \frac{6\zeta'(2)}{\pi^2} \right], \end{aligned} \quad (2)$$

where $\gamma_E \approx 0.5772$ and $\zeta'(2) \approx -0.9376$.

During the propagation of the hard parton after its production, it will suffer multiple scattering with targets in the medium. Recently many theoretical studies [3–7] have investigated the rescattering-induced radiative energy loss. Here we will focus on the stimulated emission and thermal absorption associated with rescattering in a hot QGP medium.

Assume a hard parton is produced at $\mathbf{y}_0 = (y_0, \mathbf{y}_{0\perp})$ inside the medium with y_0 being the longitudinal coordinate. The interaction between the jet and target partons is modeled by a static color-screened Yukawa potential which is initially proposed by Gyulassy-Wang (GW) [2],

$$V_n = 2\pi\delta(q^0)v(\mathbf{q}_n)e^{-i\mathbf{q}_n \cdot \mathbf{y}_n}T_{a_n}(j)T_{a_n}(n), \quad v(\mathbf{q}_n) = \frac{4\pi\alpha_s}{\mathbf{q}_n^2 + \mu^2}. \quad (3)$$

Here $\mathbf{q}_n$ is the momentum transfer from a target parton n at $\mathbf{y}_n = (y_n, \mathbf{y}_{n\perp})$, $T_{a_n}(j)$ and $T_{a_n}(n)$ are the color matrices for the jet and target parton.

Following the framework of opacity expansion developed by Gyulassy, Lévai and Vitev (GLV)[5] and Wiedemann [7], we will only consider the contributions to the first order in the opacity expansion. It was shown by GLV that the higher order corrections contribute little to the radiative energy loss. Including stimulated emission and thermal absorption we can express the radiation probability at the first order in opacity as

$$\begin{aligned} \frac{dP^{(1)}}{d\omega} &= \frac{C_2}{8\pi d_A d_R} \int \frac{dz}{z} \int \frac{d^2\mathbf{k}_\perp}{(2\pi)^2} \int \frac{d^2\mathbf{q}_\perp}{(2\pi)^2} v^2(\mathbf{q}_\perp) \frac{N}{A_\perp} \langle \text{Tr} [|R^{(S)}|^2 + 2\text{Re} (R^{(0)\dagger} R^{(D)})] \rangle \\ &\quad \times P\left(\frac{\omega}{E}\right) \left[(1 + N_g(zE)) \delta(\omega - zE) \theta(1 - z) + N_g(zE) \delta(\omega + zE) \right], \end{aligned} \quad (4)$$

where C_2 and C_A are Casimirs of the target parton in the fundamental representations in d_R and d_A dimension. Averaging over the longitudinal target profile is defined as

$\langle \dots \rangle = \int dy \rho(y) \dots$ with $\rho(y) = 2 \exp(-2y/L)/L$. We denote $R^{(0)}$, $R^{(S)}$ and $R^{(D)}$ as the radiation amplitude associated with zero, single and double Born rescattering with parton targets in the QGP, respectively. They can be obtained as[5,8]

$$R^{(0)} = 2igT_c \frac{\mathbf{k}_\perp \cdot \epsilon_\perp}{\mathbf{k}_\perp^2}, \quad (5)$$

$$R^{(S)} = 2ig(\mathbf{H}T_aT_c + \mathbf{B}_1e^{i\omega_0y_{10}}[T_c, T_a] + \mathbf{C}_1e^{i(\omega_0-\omega_1)y_{10}}[T_c, T_a]) \cdot \epsilon_\perp, \quad (6)$$

$$R^{(D)} = 2igT_ce^{i\omega_0y_{10}}\left(-\frac{C_F+C_A}{2}\mathbf{H}e^{-i\omega_0y_{10}} + \frac{C_A}{2}\mathbf{B}_1 + \frac{C_A}{2}\mathbf{C}_1e^{-i\omega_1y_{10}}\right) \cdot \epsilon_\perp. \quad (7)$$

Here $\epsilon_\perp$ is the transverse polarization of the radiative gluon, $y_{10} = y_1 - y_0$,

$$\omega_0 = \frac{\mathbf{k}_\perp^2}{2\omega}, \quad \omega_1 = \frac{(\mathbf{k}_\perp - \mathbf{q}_\perp)^2}{2\omega}, \quad (8)$$

$$\mathbf{H} = \frac{\mathbf{k}_\perp}{\mathbf{k}_\perp^2}, \quad \mathbf{C}_1 = \frac{\mathbf{k}_\perp - \mathbf{q}_\perp}{(\mathbf{k}_\perp - \mathbf{q}_\perp)^2}, \quad \mathbf{B}_1 = \mathbf{H} - \mathbf{C}_1. \quad (9)$$

The zero-temperature part of the parton energy loss corresponds to the radiation induced by rescattering which should be the same as obtained in the previous studies [5]. We denote this part as $\Delta E_{rad}^{(1)}$. The remainder or temperature-dependent part of energy loss induced by rescattering at the first order in opacity is

$$\begin{aligned} \Delta E_{abs}^{(1)} &= \int d\omega \omega \left(\frac{dP^{(1)}}{d\omega} - \frac{dP^{(1)}}{d\omega} \Big|_{T=0} \right) \\ &= \frac{\alpha_s C_F}{\pi} \frac{L}{\lambda_g} E \int dz \int \frac{d\mathbf{k}_\perp^2}{\mathbf{k}_\perp^2} \int d^2\mathbf{q}_\perp |\bar{v}(\mathbf{q}_\perp)|^2 \frac{\mathbf{k}_\perp \cdot \mathbf{q}_\perp}{(\mathbf{k}_\perp - \mathbf{q}_\perp)^2} N_g(zE) \\ &\quad \times \left[-P(-z) \langle Re(1 - e^{i\omega_1y_{10}}) \rangle + P(z) \langle Re(1 - e^{i\omega_1y_{10}}) \rangle \theta(1-z) \right], \end{aligned} \quad (10)$$

where $\lambda_g = C_F\lambda/C_A$ is the mean-free-path of the gluon, the factor $1 - \exp(i\omega_1y_{10})$ reflects the destructive interference arising from the non-Abelian Landau-Pomeranchuk-Migdal (LPM) [9] effect, and $|\bar{v}(\mathbf{q}_\perp)|^2$ is the normalized distribution of momentum transfer from the scattering centers,

$$|\bar{v}(\mathbf{q}_\perp)|^2 = \frac{1}{\pi} \frac{\mu_{eff}^2}{(\mathbf{q}_\perp^2 + \mu^2)^2}, \quad \frac{1}{\mu_{eff}^2} = \frac{1}{\mu^2} - \frac{1}{q_{\perp max}^2 + \mu^2}, \quad q_{\perp max}^2 \approx 3E\mu. \quad (11)$$

In the limit of $EL \gg 1$ and $E \gg \mu$, One can then get the approximate asymptotic behavior of the energy loss,

$$\frac{\Delta E_{rad}^{(1)}}{E} \approx \frac{\alpha_s C_F \mu^2 L^2}{4\lambda_g E} \left[\ln \frac{2E}{\mu^2 L} - 0.048 \right], \quad (12)$$

$$\frac{\Delta E_{abs}^{(1)}}{E} \approx -\frac{\pi\alpha_s C_F}{3} \frac{LT^2}{\lambda_g E^2} \left[\ln \frac{\mu^2 L}{T} - 1 + \gamma_E - \frac{6\zeta'(2)}{\pi^2} \right]. \quad (13)$$

Similarly to the final-state absorption, the minus sign in Eq. (13) indicates that the contribution from thermal absorption associated with rescattering is larger than that of stimulated emission, resulting in a net energy gain.




Figure 1. The ratio of effective parton energy loss with ($\Delta E = \Delta E_{abs}^{(0)} + \Delta E_{abs}^{(1)} + \Delta E_{rad}^{(1)}$) and without ($\Delta E_{rad}^{(1)}$) absorption as a function of E/μ . Inset box: energy gain via absorption with ($-\Delta E_{abs}^{(1)}$) and without ($-\Delta E_{abs}^{(0)}$) rescattering.

3. NUMERICAL RESULTS AND CONCLUSION

The significance of the thermal absorption relative to the induced radiation can be studied by evaluating Eqs. (2) and (10) numerically. From the perturbative QCD at finite temperature the Debye screening mass can be taken as $\mu^2 = 4\pi\alpha_s T^2$ [10]. In the GW model the mean-free-path for a gluon λ_g is [2]

$$\lambda_g^{-1} = \langle \sigma_{qg} \rho_q \rangle + \langle \sigma_{gg} \rho_g \rangle \approx \frac{2\pi\alpha_s^2}{\mu^2} 9 \times 7\zeta(3) \frac{T^3}{\pi^2}, \quad (14)$$

where $\zeta(3) \approx 1.202$. With fixed values of L/λ_g and α_s , $\Delta E/\mu$ is a function of E/μ only.

The ratios of the calculated radiative energy loss with and without stimulated emission and thermal absorption as functions of E/μ are illustrated in Fig. 1 for $L/\lambda_g = 3, 5$ and $\alpha_s = 0.3$. The thermal absorption reduces the effective parton energy loss by about 30-10% for intermediate values of parton energy. The energy dependence of the effective parton energy loss becomes stronger. This will affect the shape of suppression of p_T hadrons spectrum in the intermediate energy region. For partons with very high energy the effect of the gluon absorption is small and can be neglected.

REFERENCES

1. X.-N. Wang and M. Gyulassy, Phys. Rev. Lett. 68 (1992) 1480.
2. M. Gyulassy and X.-N. Wang, Nucl. Phys. B420 (1994) 583; X.-N. Wang, M. Gyulassy and M. Plumer, Phys. Rev. D 51 (1995) 3436.
3. R. Baier, Y. L. Dokshitzer, A. H. Mueller, S. Peigne and D. Schiff, Nucl. Phys. B 484 (1997) 265.
4. B. G. Zakharov, JETP Lett. 63 (1996) 952; JETP Lett. 65 (1997) 615.
5. M. Gyulassy, P. Levai and I. Vitev, Phys. Rev. Lett. 85 (2000) 5535.
6. X. Guo and X.-N. Wang, Phys. Rev. Lett. 85 (2000) 3591.
7. U. A. Wiedemann, Nucl. Phys. B588 (2000) 303.
8. E. Wang and X.-N. Wang, Phys. Rev. Lett. 87 (2001) 142301.
9. L. D. Landau and I. Pomeranchuk, Dokl. Akad. Nauk Ser. Fiz. 92 (1953) 735; A. B. Migdal, Phys. Rev. 103 (1956) 1811.
10. E. Braaten and R. D. Pisarski, Nucl. Phys. B337 (1990) 569.